Calculus of multivariable functions

1^{st} November 2005

Definition 1 talks about functions of n independent variables, and Definition 2 the partial derivatives of these. In Theorem 1 we have the product rule, in Theorem 2 the quotient rule, and in Theorem 3 the generalised power function rule.

Definition 1. A function $y = \mathrm{f}(x_1, \ldots, x_n)$ is called a *function of n independent variables* if there exists one and only one value of y in the range of f for each tuple of real number $(x_1, \ldots, x_n)$ in the domain of f. Here y is called the *dependent variable* while x_i, $i = 1, \ldots, n$, the *independent variables*.

§

The word *tuple* in Definition 1 means an *ordered list*. It is also known as n-tuple, where n is the size of the list.

Definition 2. Let a multivariable function be $y = \mathrm{f}(x_1, \ldots, x_n)$. The *partial derivative* of y with respect to x_i, where $1 \le i \le n$, is a measure of the instantaneous rate of change of y with respect to x_i while x_j is held constant for all $j \ne i$, where $1 \le j \le n$. This partial derivative is defined as

$$\frac{\partial y}{\partial x_i} = \lim_{\Delta x_i \to 0} \frac{\mathrm{f}(\ldots, x_i + \Delta x_i, \ldots) - \mathrm{f}(x_1, \ldots, x_n)}{\Delta x_i}$$

and can be written in either one of the following forms.

$$\frac{\partial y}{\partial x_i},\ \frac{\partial \mathrm{f}}{\partial x_i},\ \mathrm{f}_{x_i}(x_1, \ldots, x_n),\ \mathrm{f}_{x_i},\ \text{or}\ y_{x_i}$$

§

Theorem 1. Let $z = \mathrm{g}(x, y) \cdot \mathrm{h}(x, y)$. Then,

$$\frac{\partial z}{\partial x} = \mathrm{g} \cdot \frac{\partial \mathrm{h}}{\partial x} + \mathrm{h} \cdot \frac{\partial \mathrm{g}}{\partial x}$$

and

$$\frac{\partial z}{\partial y} = \mathrm{g} \cdot \frac{\partial \mathrm{h}}{\partial y} + \mathrm{h} \cdot \frac{\partial \mathrm{g}}{\partial y}$$

§

Theorem 2. Let $z = \frac{\mathrm{g}(x,y)}{\mathrm{h}(x,y)}$ and $\mathrm{h}(x, y) \ne 0$. Then,

$$\frac{\partial z}{\partial x} = \frac{\mathrm{h} \cdot \frac{\partial \mathrm{g}}{\partial x} - \mathrm{g} \cdot \frac{\partial \mathrm{h}}{\partial x}}{\mathrm{h}^2}$$

and

$$\frac{\partial z}{\partial y} = \frac{\mathrm{h} \cdot \frac{\partial \mathrm{g}}{\partial y} - \mathrm{g} \cdot \frac{\partial \mathrm{h}}{\partial y}}{\mathrm{h}^2}$$

§

Theorem 3. Let $z = [\mathrm{g}(x, y)]^n$. Then,

$$\frac{\partial z}{\partial x} = n\mathrm{g}^{n-1} \cdot \frac{\partial \mathrm{g}}{\partial x}$$

and

$$\frac{\partial z}{\partial y} = n\mathrm{g}^{n-1} \cdot \frac{\partial \mathrm{g}}{\partial y}$$

§

Exercise 1. Find the first-order partial derivatives of the following:

1. $c = 100(1 + e^{-1.2q})$	*2.* $q = 5l^{0.7}k^{0.3} - 4l - 3k + 10$	*3.* $q = \ln 3x + 2x \ln y$
4. $z = x^3(1 + y + y^2)$	*5.* $p = 150e^{0.74t}$	*6.* $q = lnx + \ln y$
7. $z = x^3 + x^2 + x + 2xy + xy^2$	*8.* $z = x^2y^5$	*9.* $q = 2l^{0.77}k^{0.1}$
10. $u = 8x^3y^3$	*11.* $z = xy + \frac{y}{x}$	*12.* $u = 7l^7k^4$

§

In Definition 3 we find the meaning of second-order partial derivatives. Theorem 4 is about critical points, and Procedure 1 is a procedure for determining critical points.

Definition 3. Let $z = \mathrm{f}(x, y)$. Then, the *second-order direct partial derivatives* are

$$\frac{\partial}{\partial x}\left(\frac{\partial z}{\partial x}\right) \quad \text{and} \quad \frac{\partial}{\partial y}\left(\frac{\partial z}{\partial y}\right)$$

These are also written

$$\mathrm{f}_{xx},\ (\mathrm{f}_x)_x,\ \frac{\partial^2 z}{\partial x^2} \quad \text{and respectively} \quad \mathrm{f}_{yy},\ (\mathrm{f}_y)_y,\ \frac{\partial^2 z}{\partial y^2}$$

The *cross partial derivatives* are

$$\frac{\partial}{\partial y}\left(\frac{\partial z}{\partial x}\right) \quad \text{and} \quad \frac{\partial}{\partial x}\left(\frac{\partial z}{\partial y}\right)$$

These are also written as

$$\mathrm{f}_{xy},\ (\mathrm{f}_x)_y,\ \frac{\partial^2 z}{\partial y \partial x} \quad \text{and respectively} \quad \mathrm{f}_{yx},\ (\mathrm{f}_y)_x,\ \frac{\partial^2 z}{\partial x \partial y}$$

§

Exercise 2. Find the second-order partial derivatives of the following:

1. $z = x^3 + xy$	*2.* $z = l^{0.2}k^{0.3} - \lambda(10 - l - 3k)$	*3.* $z = 2x^3y^4$
4. $c = 100(1 + e^{-0.1q})$	*5.* $q = \ln x + \ln y$	*6.* $q = 5l^{0.44}k^{0.2} + 50 - 7l - 5k$
7. $q = 9l^{0.77}k^{0.1}$	*8.* $q = \ln 5x + x \ln y$	*9.* $p = 110e^{0.87t}$
10. $z = xy + \frac{y}{2x}$	*11.* $u = 10x^4y^4$	*12.* $z = x^3(1 + y + y^3)$

§

When the second derivative is negative, the curve is concave towards the origin.

Theorem 4. For a multivariable function $z = \mathrm{f}(x, y)$ to be a *relative maximum* at (a, b) necessarily $\mathrm{f}_x, \mathrm{f}_y = 0$, and $\mathrm{f}_{xx}, \mathrm{f}_{yy} < 0$ and $\mathrm{f}_{xx} \cdot \mathrm{f}_{yy} > (\mathrm{f}_{xy})^2$ at that point. For the same at the same to be a *relative minimum*, necessarily $\mathrm{f}_x, \mathrm{f}_y = 0$, and $\mathrm{f}_{xx}, \mathrm{f}_{yy} > 0$ and $\mathrm{f}_{xx} \cdot \mathrm{f}_{yy} > (\mathrm{f}_{xy})^2$ there. Moreover, an *inflection point* is a point (a, b) at which $\mathrm{f}_{xx} \cdot \mathrm{f}_{yy} < (\mathrm{f}_{xy})^2$, and both f_{xx} and f_{yy} have the same sign. On the other hand, a *saddle point* is a point (a, b) at which $\mathrm{f}_{xx} \cdot \mathrm{f}_{yy} < (\mathrm{f}_{xy})^2$, but f_{xx} and f_{yy} are of different signs.

§

Procedure 1 *Procedure for determining a critical point of a function with two independent variables*

Given $z = \mathrm{f}(x, y)$ and a point (a, b)**, at this point,**
if $\mathrm{f}_x = 0$ and $\mathrm{f}_y = 0$ **then**
  (a, b) is a critical point
  if $\mathrm{f}_{xx} \cdot \mathrm{f}_{yy} > (\mathrm{f}_{xy})^2$ **then**
    if $\mathrm{f}_{xx} < 0$ and $\mathrm{f}_{yy} < 0$ **then**
      (a, b) is a relative maximum of z
    elseif $\mathrm{f}_{xx} > 0$ and $\mathrm{f}_{yy} > 0$ **then**
      (a, b) is a relative minimum of z
    else †
    endif
  elseif $\mathrm{f}_{xx} \cdot \mathrm{f}_{yy} < (\mathrm{f}_{xy})^2$ **then**
    if $\mathrm{f}_{xx} \cdot \mathrm{f}yy > 0$ **then**
      (a, b) is an inflection point
    elseif $\mathrm{f}_{xx} \cdot \mathrm{f}_{yy} < 0$ **then**
      (a, b) is a saddle point
    else ‡
    endif
  else
    test inconclusive
  endif
else
  (a, b) is no critical point
endif

§

Problem 1. There are two dead ends in Procedure 1. The first one (†) is the case where $\mathrm{f}_{xx} \cdot \mathrm{f}_{yy} > (\mathrm{f}_{xy})^2$ and either $(\mathrm{f}_{xx} = 0, \mathrm{f}_{yy} = 0)$, $(\mathrm{f}_{xx} = 0, \mathrm{f}_{yy} < 0)$, $(\mathrm{f}_{xx} = 0, \mathrm{f}_{yy} > 0)$, $(\mathrm{f}_{xx} < 0, \mathrm{f}_{yy} = 0)$, $(\mathrm{f}_{xx} > 0, \mathrm{f}_{yy} = 0)$, $(\mathrm{f}_{xx} < 0, \mathrm{f}_{yy} > 0)$, or $(\mathrm{f}_{xx} > 0, \mathrm{f}_{yy} < 0)$. The second one (‡) is where $\mathrm{f}_{xx} \cdot \mathrm{f}_{yy} = 0$. Find out what happen in these cases, and thus complete the missing lines of logic in Procedure 1.

§

Definition 4 gives the meaning of derivative and differential. Example 1 looks at the derivative and differential of functions of one variable and two variables.

Definition 4. By *derivative* $\frac{\mathrm{d}y}{\mathrm{d}x}$ we mean an infinitesimally small change in y with respect to an infinitesimally small change in x. By *differential* $\mathrm{d}y$ and $\mathrm{d}x$ we mean an infinitesimally small change in the values of y and respectively x.

§

Example 1. For a function of one variable $y = \mathrm{f}(x)$, the total derivative is

$$\frac{\mathrm{d}y}{\mathrm{d}x}$$

and the differential of y is

$$\mathrm{d}y = \left(\frac{\mathrm{d}y}{\mathrm{d}x}\right)\mathrm{d}x$$

For a function of two variables $z = \mathrm{f}(x, y)$ partial derivatives are, the first-order partial derivatives

$$\frac{\partial z}{\partial x} \quad \text{and} \quad \frac{\partial z}{\partial y}$$

and the second-order partial derivatives

$$\frac{\partial^2 z}{\partial x^2} \equiv z_{xx},\ \frac{\partial^2 z}{\partial y^2} \equiv z_{yy},\ \frac{\partial^2}{\partial y \partial x} \equiv z_{xy} \text{ and } \frac{\partial^2 z}{\partial x \partial y} \equiv z_{yx}$$

The total differential of z is

$$\mathrm{d}z = \left(\frac{\partial \mathrm{f}}{\partial x}\right)\mathrm{d}x + \left(\frac{\partial \mathrm{f}}{\partial y}\right)\mathrm{d}y$$

and for small changes which are not infinitesimal, $\mathrm{d}x$ becomes Δx and the incremental change formula is

$$\Delta z \approx \left(\frac{\partial \mathrm{f}}{\partial x}\right)\Delta x + \left(\frac{\partial \mathrm{f}}{\partial y}\right)\Delta y$$

Definition 5 gives the general production function. In Example 2 we look at the Cobb-Douglas production function in more details. Theorem 5 gives the law of diminishing returns to labour and the proof thereof, while similarly does Theorem 6 the law of diminishing returns to capital.

Definition 5. The *general production function* is $q = \mathrm{f}(l, k)$, where q is output of the production, l labour and k capital. The *Cobb-Douglas production function* in its general form is

$$q = a l^{\alpha} k^{\beta} \tag{1}$$

where a is a constant and $0 < \alpha < 1$, $0 < \beta < 1$, $l > 0$ and $k > 0$.

§

Example 2. With the Cobb-Douglas production function, the *marginal product of labour* is,

$$p_{lm} = q_l = \frac{\partial q}{\partial l} = a\alpha l^{\alpha-1} k^{\beta} \tag{2}$$

and the *marginal product of capital*

$$p_{km} = q_k = \frac{\partial q}{\partial k} = a\beta l^{\alpha} k^{\beta-1} \tag{3}$$

From this we see that $p_{lm} > 0$ and $p_{km} > 0$.

Theorem 5. From the Cobb-Douglas production function we have the *law of diminishing returns to labour*, which states that $q_{ll} < 0$.

Proof. From Equation 1 in Definition 5,

$$q_{ll} = \frac{\partial^2 q}{\partial l^2} = \frac{\partial}{\partial l}\left(\frac{\partial q}{\partial l}\right) = \frac{\partial p_{lm}}{\partial l} = (\alpha - 1)\frac{\alpha q}{l^2}$$

Since $0 < \alpha < 1$, therefore $q_{ll} < 0$. ¶

Theorem 6. Using the Cobb-Douglas production function, the *law of diminishing returns to capital* states that $q_{kk} < 0$.

Proof. From Equation 1 in Definition 5,

$$q_{kk} = \frac{\partial^2 q}{\partial k^2} = \frac{\partial}{\partial k}\left(\frac{\partial q}{\partial k}\right) = \frac{\partial p_{km}}{\partial k} = (\beta - 1)\frac{\beta q}{k^2}$$

which, with $0 < \beta < 1$, tells us that $q_{kk} < 0$. ¶

In Example 3 we see the changes in marginal product values.

Example 3. Using the Cobb-Douglas production function,

$$q_{kl} = q_{lk} = a\alpha\beta l^{\alpha-1}k^{\beta-1}$$

Therefore, $q_{lk} > 0$ and $q_{kl} > 0$. In other words, p_{lm} increases as capital input k increases, and respectively p_{km} increases as labour input l increases.

Example 4 shows us the average functions of labour and capital, Example 5 the marginal functions of labour and capital, and Example 6 the comparison between marginal and average functions.

Example 4. For the Cobb-Douglas production function in Equation 1 the average product of labour is

$$p_{la} = \frac{q}{l} = al^{\alpha-1}k^{\beta} \tag{4}$$

and the average product of capital is

$$p_{ka} = \frac{q}{k} = al^{\alpha}k^{\beta-1} \tag{5}$$

Example 5. Again using the Cobb-Douglas production function of Equation 1, the marginal product of labour is

$$p_{lm} = \frac{\partial q}{\partial l} = a\alpha l^{\alpha-1}k^{\beta} \tag{6}$$

and the marginal product of capital is

$$p_{km} = \frac{\partial q}{\partial k} = a\beta l^{\alpha}k^{\beta-1} \tag{7}$$

Example 6. From the APL equation, Equation 4, and the MPL equation, Equation 6, and since $0 < \alpha < 1$, therefore $p_{ml} < p_{la}$. Similarly from the APK equation, Equation 5, and the MPK equation, Equation 7, since $0 < \beta < 1$, we have $p_{km} < p_{ka}$.

In Example 7 one sees the conditions for using labour, Equation 8, and the conditions for using capital, Equation 9.

Example 7. A producer likes to have a positive marginal function, which means that the productivity increases as the input increases. But the second derivative is negative, which means that this rate of increase slows down as time goes by. In practice, the conditions for using labour are,

$$p_{lm} = \frac{\partial q}{\partial l} > 0,\ \frac{\mathrm{d}p_{lm}}{\mathrm{d}l} = \frac{\partial^2 q}{\partial l^2} < 0, \text{ and } p_{lm} < p_{la} \tag{8}$$

The conditions for using capital are similarly,

$$p_{km} = \frac{\partial q}{\partial k} > 0,\ \frac{\mathrm{d}p_{km}}{\mathrm{d}k} = \frac{\partial^2 q}{\partial k^2} < 0, \text{ and } p_{km} < p_{ka} \tag{9}$$

Definition 6 and Theorem 7 deal respectively with production function graphs and slope of an isoquant.

Definition 6. An *isoquant* is a graph in two dimensions, $k = \mathrm{f}(l)$, plotted to represent a production function $q = \mathrm{f}(l, k)$. The slope $\frac{\mathrm{d}k}{\mathrm{d}l}$ is called the *marginal rate of technical substitution*. The value of this slope at (l_0, k_0) is denoted by $\frac{\mathrm{d}k}{\mathrm{d}l}\big|_{l_0 k_0}$.

§

Theorem 7. The slope of an isoquant is the ratio of the marginal products.

Proof. The total differential of $q = \mathrm{f}(l, k)$ is

$$\mathrm{d}q = \left(\frac{\partial q}{\partial l}\right)\mathrm{d}l + \left(\frac{\partial q}{\partial k}\right)\mathrm{d}k$$

Along any isoquant, $\mathrm{d}q = 0$, therefore,

$$0 = \left(\frac{\partial q}{\partial l}\right)\mathrm{d}l + \left(\frac{\partial q}{\partial k}\right)\mathrm{d}k \tag{10}$$

This directly yield, after some manipulation,

$$\frac{\mathrm{d}k}{\mathrm{d}l} = -\frac{q_l}{q_k}$$

Or, from Equation 10 together with Equation's 6 and 7, it follows that,

$$\frac{\mathrm{d}k}{\mathrm{d}l} = -\frac{p_{lm}}{p_{km}}$$

Definition 7. In the Cobb-Douglas production function equation, Equation 1, let both inputs l and k change by the same proportion, and let λ be the constant of this proportionality. Then $q_2 = a(\lambda l)^{\alpha}(\lambda k)^{\beta}$, which leads to $q_2 = \lambda^{\alpha+\beta} q_1$. When $\alpha + \beta = 1$, the case is described as *constant returns to scale*, when $\alpha + \beta < 1$ as *decreasing returns to scale*, and when $\alpha + \beta > 1$ as *increasing returns to scale*.

§

Definition 8. A *homogeneous* Cobb-Douglas production function of order r is,

$$\mathrm{f}(\lambda l, \lambda k) = \lambda^r \mathrm{f}(l, k)$$

where $r = (\alpha, \beta)$.

§

The utility function in its general form is given in Definition 9. Definition 10 gives the Cobb-Douglas utility function, Definition 11 the marginal utility, and Definition 12 the meaning of indifference curves.

Definition 9. A *utility function* expresses utility as a function of goods consumed. In its general form this is,

$$u = \mathrm{f}(x, y)$$

where x and y are the quantities of goods X and respectively Y consumed.

§

Definition 10. The *Cobb-Douglas utility function* is in its general form,

$$u = ax^{\alpha} y^{\beta}$$

where a is a constant, and $0 < \alpha < 1$, $0 < \beta < 1$, $x > 0$ and $y > 0$.

§

Definition 11. The *marginal utility* for a utility function with one variable, $u = \mathrm{f}(x)$, is $\frac{\mathrm{d}u}{\mathrm{d}x} = u_x = u_{xm}$. The marginal utility for a utility function with two variables, $u = \mathrm{f}(x, y)$, is $\frac{\partial u}{\partial x} = u_x = u_{xm}$ and $\frac{\partial u}{\partial y} = u_y = u_{ym}$.

§

Definition 12. The *indifference curve* is a graph $y = \mathrm{f}(x)$ drawn to represent a utility function $u = \mathrm{f}(x, y)$. Its slope $\frac{\mathrm{d}y}{\mathrm{d}x}$ is called the *marginal rate of substitution*. Setting the total differential equal to zero,

$$0 = \mathrm{d}u = \left(\frac{\partial u}{\partial x}\right)\mathrm{d}x + \left(\frac{\partial u}{\partial y}\right)\mathrm{d}y$$

we find

$$\frac{\mathrm{d}y}{\mathrm{d}x} = -\frac{u_x}{u_y}$$

and

$$\frac{\mathrm{d}y}{\mathrm{d}x} = -\frac{u_{xm}}{u_{ym}}$$

§

Partial elasticities are describe in Definition 13, partial elasticities of demand, and Example's 8 and 9, respectively the partial elasticity with respect to labour and the partial elasticity with respect to capital.

Definition 13. Let a demand function be

$$q_a = \mathrm{f}(p_a, y, p_b) \tag{11}$$

where q_a is the quantity demanded of good a, p_a the price of a, y consumer's income, and p_b the price of another good b. Then, the *price elasticity of demand* is,

$$\varepsilon_d = \frac{\partial q_a}{\partial p_a}\frac{p_a}{q_a}$$

The *income elasticity of demand* is,

$$\varepsilon_y = \frac{\partial q_a}{\partial y}\frac{y}{q_a}$$

And the *cross-price elasticity of demand* is,

$$\varepsilon_c = \frac{\partial q_a}{\partial p_b}\frac{p_b}{q_a}$$

§

Example 8. With the demand function as in Equation 11, the partial elasticity with respect to labour is,

$$\varepsilon_{ql} = \frac{\partial q}{\partial l}\frac{l}{q}$$

And from Equation's 6 and 4, this leads to,

$$\varepsilon_{ql} = \frac{p_{lm}}{p_{la}}$$

For the Cobb-Douglas production function, Equation 1, then $\varepsilon_{ql} = \alpha$.

§

Example 9. Again, with the demand function as in Equation 11, the partial elasticity with respect to capital is,

$$\varepsilon_{qk} = \frac{\partial q}{\partial k}\frac{k}{q}$$

Then, from Equation's 7 and 5,

$$\varepsilon_{qk} = \frac{p_{km}}{p_{ka}}$$

For the Cobb-Douglas production function, Equation 1, we have $\varepsilon_{qk} = \beta$.

§

Bibliography

Edward T Dowling. *Introduction to mathematical economics.* Schaum's outline series,2nd ed. 1992(1980)

Teresa Bradley. *Essential mathematics for economics and business.* 2nd ed. 2002